








10,200 Treated Injuries

 2007-2022 There has been an upward trend of 535 injuries seen in the ED in the US per year

- **7,400** Injuries were during a 1-month period between June 17, 2022- July 17, 2022
 - 65% of those injuries were Male
 - **1,300** of these injuries were firecrackers
 - 600 were from **SPARKLERS**
- Ages 25-44 experienced 36% of the injuries
 - Children <15 experienced 28%
 - >65 accounted for 3%
- 11 DEATHS










BURNS First Degree


Maintained at home or See your PCP

- Redness that blanches, swelling, mild pain
- Cleanse the area gently with soap and tepid water, cool compress, moisturize with aloe or after-sun cream





CAN WE TREAT THIS AT HOME?






BURNS

Second Degree

Seek medical attention if >3in in diameter, or in a sensitive area (hands, feet, face, genitals, buttocks, or over a major joint)

- Blisters, red/pink, may or may not blanch depending on severity, wet and weeping with serous fluid
- Cool water, NO ICE, apply a cool compress for about 30 min, antibiotic cream (OTC or RX) no stick bandage and wrap loosely, elevate to decrease swelling (change bandage 3xday)










BURNS

Third Degree

SEEK IMMEDIATE MEDICAL ATTENTION

- Wax-white to leather grey to charred black, no blanching, dry, firm, absence of hair can extend into the fascia/muscle
- Often requires skin grafting and possibly amputation











CAN WE TREAT THIS AT HOME?





Resources

- 2022 Annual Fireworks Report (cpsc.gov)
- Fireworks | CPSC.gov